

## Northeast Dental Group

'A Dental Destination Location'

7770 Frontage Rd #7341 Cicero, NY 13039

## **CONSENT FOR ALVEOLOPLASTY**

Date

Patient's Name

	nitial each paragraph after reading. If you have any questions, please ask tor BEFORE initialing.
	the right to be informed about your diagnosis and planned surgery so that you le whether to have a procedure or not after knowing the risks and benefits.
Your plan Alternativ Taking te	nosis is: bone spicule (sharp bone) ned treatment is:alveoloplasty with Local Anesthesia e treatment methods include:No Treatment/Partial Treatment/Defer Treatment. eth out is a permanent process. Whether the procedure is easy or difficult, it is gical procedure. All surgeries have some risks. They include the following and Swelling, bruising and pain.
2.	Stretching of the corners of the mouth that may lead to cracking or bruising.
3.	Possible infection that might need more treatment.
4.	Dry socket - jaw pain that begins a few days after surgery, that may need more care.
5.	Possible damage to other teeth close to the ones being taken out, more often those with large fillings or caps.
6.	Numbness, pain, or changed feelings in the teeth, gums, lip, chin and/or tongue (including possible loss of taste). This is due to the closeness of tooth roots (mainly with wisdom teeth) to the nerves which can be injured or damaged. Usually the numbness or pain goes away, but in some cases, it may need more treatment or may be permanent.
7.	Trismus – you can only open your mouth a little. This is most common after wisdom teeth are taken out. Sometimes it happens because of jaw joint (TMJ) problems already there. Damage can occur to the ligaments of the jaw joint (TMJ) from having your mouth open wide and/or for a period of time. This is more common if you already have symptoms or signs. This may need separate additional treatment.

8.	Bleeding – oozing can often happen for sev not common.	eral hours, but a lot of bleeding is
9.	Sharp ridges or bone splinters may form la the tooth was taken out. These may nee remove.	
10.	Sometimes tooth roots may be left in to avo	• .
11.	The roots of the upper back teeth are often a piece of root can get into the sinus. An of into the mouth that may need more treatmen	ppening may occur from the sinus
12.	It is very rare that the jaw will break, but it is are buried very deep in their sockets.	possible in cases where the teeth
13.	When donated, processed, or artificial bone a socket the pieces might not join together volst.	•
14.	It is understood all encounters at Northe consultation/surgery/follow-up/phone calls me training and/or documentation. This recompermanent dental record or may be discardental office.	nay be recorded for the purpose of ording may become part of my
	I have told my doctor that I use birth control some antibiotics and other medications made birth control pills, and I could conceive and discuss with my personal doctor using other treatment, and to continue those methods used and use only oral birth control.	ry reduce the preventive effect of ad become pregnant. I agree to be forms of birth control during my ntil my personal doctor says that I
CONSEN		
that the to choices I and unde have give pregnance	reatment listed above and other forms of treetment listed above and other forms of treethave and the risks of those choices have be restand the above and give my consent to seen a complete and truthful medical history, it y, etc. I certify that I speak, read and write I wered before signing this form.	eatment or no treatment at all are een presented to me. I have read urgery and chosen anesthesia. I ncluding all medicines, drug use,
Detient's	(or Logal Cuardian's) Signature	Data
rauents (	(or Legal Guardian's) Signature	Date
Doctor's S	Signature	Date

Date

Witness' Signature